



ELSEVIER

Journal of Pure and Applied Algebra 123 (1998) 51–66

JOURNAL OF
PURE AND
APPLIED ALGEBRA

Rings whose free modules satisfy the ascending chain condition on submodules with a bounded number of generators

Maria Elisa Antunes Simões^a, Patrick F. Smith^{b,*}

^a *Departamento de Matemática, Faculdade de Ciências, Universidade de Lisboa, Lisbon, Portugal*

^b *Department of Mathematics, University of Glasgow, Glasgow G12 8QW, UK*

Communicated by C.A. Weibel; received 6 March 1995; revised 19 March 1996

Abstract

Let R be a ring such that every finitely generated free (respectively, every free) right R -module satisfies the ascending chain condition on n -generated submodules for every positive integer n ; then any ring Morita equivalent to R has the same property. This is in contrast to rings R which satisfy the ascending chain condition on n -generated right ideals, for some fixed positive integer n , for in this case rings Morita equivalent to R need not have the same property. If R is a right and left Ore domain and n is a positive integer such that the free right R -module $R_R^{(n)}$ satisfies the ascending chain condition on n -generated submodules then so too does every free right R -module. Many examples are given of rings for which every finitely generated free (respectively, every free) right module satisfies the ascending chain condition on n -generated submodules, for some positive integer n . © 1998 Elsevier Science B.V.

AMS Classification: 16P70, 16D90, 16U20, 16D40, 16P20

1. Morita equivalence

Throughout this note, all rings are associative with identity and all modules are unital right modules. Let n be a positive integer. We say that a module M satisfies *n-acc* if every ascending chain of n -generated submodules terminates. If the module M satisfies *n-acc* for every positive integer n , then we shall say that M satisfies *pan-acc*. We shall say that the ring R satisfies *right n-acc* (respectively, *right pan-acc*) if the right

* Corresponding author.

R -module R satisfies n -acc (*pan-acc*). For information about any terms used without explanation, see [1] or [9].

Let R be any ring and let m, n be positive integers. Let $\mathcal{M}_n(R)$ denote the ring of all $n \times n$ matrices with entries in R and let $\mathcal{M}_{m \times n}(R)$ denote the additive Abelian group of all $m \times n$ matrices with entries in R . Let S denote the ring $\mathcal{M}_n(R)$. Clearly $\mathcal{M}_{m \times n}(R)$ is a right S -module with respect to matrix multiplication. Given elements $a_{ij} \in R$ ($1 \leq i \leq m$, $1 \leq j \leq n$), let (a_{ij}) denote the $m \times n$ matrix

$$\begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \cdots & a_{mn} \end{bmatrix}$$

in $\mathcal{M}_{m \times n}(R)$.

Let F denote the free right R -module $R_R^{(m)}$. Let N and L be any n -generated R -submodules of F . There exist $a_{ij}, b_{ij} \in R$ ($1 \leq i \leq m$, $1 \leq j \leq n$) such that

$$N = (a_{11}, \dots, a_{m1})R + \cdots + (a_{1n}, \dots, a_{mn})R,$$

and

$$L = (b_{11}, \dots, b_{m1})R + \cdots + (b_{1n}, \dots, b_{mn})R.$$

Lemma 1.1. *With the above notation, $N \subseteq L$ if and only if there exists (c_{ij}) in S such that $(a_{ij}) = (b_{ij})(c_{ij})$.*

Proof. $N \subseteq L$ if and only if there exist elements $c_{ij} \in R$ ($1 \leq i, j \leq n$) such that

$$(a_{1j}, \dots, a_{mj}) = (b_{11}, \dots, b_{m1})c_{1j} + \cdots + (b_{1n}, \dots, b_{mn})c_{nj},$$

for each $1 \leq j \leq n$, and this holds if and only if

$$\begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \cdots & a_{mn} \end{bmatrix} = \begin{bmatrix} b_{11} & \cdots & b_{1n} \\ \vdots & & \vdots \\ b_{m1} & \cdots & b_{mn} \end{bmatrix} \begin{bmatrix} c_{11} & \cdots & c_{1n} \\ \vdots & & \vdots \\ c_{n1} & \cdots & c_{nn} \end{bmatrix}. \quad \square$$

With N as above, we define $\alpha(N) = (a_{ij})S$, i.e. $\alpha(N)$ is the set of $m \times n$ matrices over R such that the transpose of each column is in N . Note that by Lemma 1.1, $\alpha(N)$ is independent of the choice of n -generating set for N . Moreover, Lemma 1.1 gives at once:

Corollary 1.2. *With the above notation, let $N \subseteq L$ be n -generated R -submodules of F . Then $\alpha(N) \subseteq \alpha(L)$.*

Now suppose that $a = (a_{ij}) \in \mathcal{M}_{m \times n}(R)$, where $a_{ij} \in R$ ($1 \leq i \leq m$, $1 \leq j \leq n$). We define

$$\beta(aS) = (a_{11}, \dots, a_{m1})R + \cdots + (a_{1n}, \dots, a_{mn})R.$$

Note that Lemma 1.1 shows that if $a, b \in \mathcal{M}_{m \times n}(R)$ with $aS = bS$ then $\beta(aS) = \beta(bS)$. Thus, for each a in $\mathcal{M}_{m \times n}(R)$, $\beta(aS)$ is a well-defined n -generated R -submodule of F . Moreover, Lemma 1.1 gives at once:

Corollary 1.3. *Let $a, b \in \mathcal{M}_{m \times n}(R)$ with $aS \subseteq bS$. Then $\beta(aS) \subseteq \beta(bS)$.*

Lemma 1.4. *With the above notation, $\beta\alpha(N) = N$ for every n -generated R -submodule N of F and $\alpha\beta(aS) = aS$ for every $a \in \mathcal{M}_{m \times n}(R)$.*

Proof. Clear. \square

Theorem 1.5. *Let R be any ring and let m and n be positive integers. Then the free right R -module $R_R^{(m)}$ satisfies n -acc if and only if the right $\mathcal{M}_n(R)$ -module $\mathcal{M}_{m \times n}(R)$ satisfies 1-acc.*

Proof. Clear by the above results. \square

Corollary 1.6. *Let R be any ring and let n be any positive integer. Then the ring $\mathcal{M}_n(R)$ satisfies right 1-acc if and only if the free right R -module $R_R^{(n)}$ satisfies n -acc.*

Proof. Take $m = n$ in the theorem. \square

Corollary 1.7. *Let R be any ring and let n be any positive integer such that the ring $\mathcal{M}_n(R)$ satisfies right 1-acc. Then R satisfies right n -acc.*

Proof. By Corollary 1.6. \square

Heinzer and Lantz [7, Section 4] show that for every positive integer n there exists a commutative ring R_n such that R_n satisfies n -acc but R_n does not satisfy $(n+1)$ -acc. Thus $\mathcal{M}_{n+1}(R_n)$ does not satisfy 1-acc (Corollary 1.7). This shows that for any positive integer n , matrix rings over rings which satisfy right n -acc need not themselves satisfy right n -acc, and in particular “satisfying right n -acc” is not a Morita invariant.

Let R be any ring and let $S = \mathcal{M}_n(R)$, for any positive integer n . For each $1 \leq i, j \leq n$, let e_{ij} denote the matrix in S with (i, j) th entry 1 and all other entries 0. Let F be a free right S -module with basis $\{f_\lambda: \lambda \in \Lambda\}$. Then F is a free right R -module with basis $\{f_\lambda e_{ij}: \lambda \in \Lambda, 1 \leq i, j \leq n\}$, and if N is any m -generated S -submodule of F , say $N = x_1 S + \cdots + x_m S$ then N is an mn^2 -generated R -submodule of F , because $N = \sum_i \sum_j \sum_k x_k e_{ij} R$. This gives the following result.

Lemma 1.8. *Let R be any ring such that every (finitely generated) free right R -module satisfies pan-acc. Let n be any positive integer. Then every (finitely generated) free right $\mathcal{M}_n(R)$ -module satisfies pan-acc.*

Theorem 1.9. *The following statements are equivalent for a ring R :*

- (i) *For each positive integer n , the ring $\mathcal{M}_n(R)$ satisfies right pan-acc.*
- (ii) *For each positive integer n , the ring $\mathcal{M}_n(R)$ satisfies right 1-acc.*
- (iii) *Every finitely generated free right R -module satisfies pan-acc.*
- (iv) *For each positive integer n , every finitely generated free right $\mathcal{M}_n(R)$ -module satisfies pan-acc.*

Proof. (i) \Rightarrow (ii): Clear.

(ii) \Rightarrow (iii): Let m be any positive integer and let $F = R_R^{(m)}$. Let n be any positive integer. By hypothesis, the ring $\mathcal{M}_{m+n}(R)$ satisfies right 1-acc. By Corollary 1.6, $R_R^{(m+n)}$ satisfies $(m+n)$ -acc. Hence F satisfies n -acc. It follows that F satisfies pan-acc. This proves (iii).

(iii) \Rightarrow (iv): By Lemma 1.8.

(iv) \Rightarrow (i): Clear. \square

Renault [11] gives an example of a right Noetherian ring R with the property that if F is the free right R -module of countably infinite rank then F does not satisfy 1-acc. Thus every finitely generated free right R -module is Noetherian and hence satisfies pan-acc but not every free right R -module satisfies pan-acc. If we assume in Theorem 1.9 that the ring R has additional properties then we can say more.

Corollary 1.10. *Let R be a right Goldie ring which satisfies dcc on right annihilators. Then the following statements are equivalent:*

- (i) *For each positive integer n , the ring $\mathcal{M}_n(R)$ satisfies right pan-acc.*
- (ii) *For each positive integer n , the ring $\mathcal{M}_n(R)$ satisfies right 1-acc.*
- (iii) *Every free right R -module satisfies pan-acc.*
- (iv) *For each positive integer n , every free right $\mathcal{M}_n(R)$ -module satisfies pan-acc.*

Proof. By Theorem 1.9 and [4, Theorem 1].

In particular, Corollary 1.10 holds for any right nonsingular right Goldie ring (see [4] or [3, Theorem 1.5]).

Lemma 1.11. *Let T be a ring, let e be an idempotent in T and let R be the subring eTe of T . Let n be any positive integer.*

- (i) *If T satisfies right n -acc then so too does R .*
- (ii) *If every (finitely generated) free right T -module satisfies n -acc then so too does every (finitely generated) free right R -module.*

Proof. (i) See [5, Proposition 4.6].

(ii) Let F be any free right R -module. Without loss of generality we can take $F = R_R^{(\Lambda)}$, for some index set Λ . We can think of F as an R -submodule of the free right T -module $G = T_T^{(\Lambda)}$, in a natural way. Let N be any n -generated R -submodule

of F , say $N = x_1R + \cdots + x_nR$. Then

$$NT = x_1RT + \cdots + x_nRT = x_1T + \cdots + x_nT \subseteq NT,$$

so that NT is an n -generated T -submodule of G .

Let $N_1 \subseteq N_2 \subseteq N_3 \subseteq \cdots$ be any ascending chain of n -generated R -submodules of F . Then, by the above remarks, $N_1T \subseteq N_2T \subseteq N_3T \subseteq \cdots$ is an ascending chain of n -generated T -submodules of G . By hypothesis, there exists a positive integer k such that $N_kT = N_{k+1}T = N_{k+2}T = \cdots$. Let $i \geq k$. Then $N_k = N_kR = N_k(eTe) = N_kTe = N_iTe = N_i$. That is, $N_k = N_{k+1} = N_{k+2} = \cdots$. It follows that F satisfies n -acc. \square

Theorem 1.12. *Let R be a ring such that every (finitely generated) free right R -module satisfies pan -acc. Let T be a ring Morita equivalent to R . Then every (finitely generated) free right T -module satisfies pan -acc.*

Proof. By Lemmas 1.8 and 1.11. \square

Let R be a ring which satisfies right pan -acc and let T be a ring Morita equivalent to R . Does T satisfy right pan -acc? By Theorem 1.12, this is certainly the case if every finitely generated free right R -module satisfies pan -acc. Heinzer and Lantz conjecture that if a ring R satisfies right pan -acc then every finitely generated free right R -module satisfies pan -acc, but this is still open according to Bonang [5] (see also [6, Ex. 0.1]). We shall return to this question in the next section.

2. Domains with n -acc

The purpose of this section is to give a proof of the main result of this paper, namely:

Theorem 2.1. *Let R be a left and right Ore domain and let n be a positive integer such that the free right R -module $R_R^{(n)}$ satisfies n -acc. Then every free right R -module satisfies n -acc.*

Combining this theorem with our remarks at the end of the previous section we see that if R is a left and right Ore domain such that for every positive integer n , the free right R -module $R_R^{(n)}$ satisfies n -acc then every ring Morita equivalent to R satisfies right pan -acc.

In order to prove Theorem 2.1 we first prove a number of lemmas.

Lemma 2.2. *Let D be a division ring and let $a \in \mathcal{M}_{m \times n}(D)$ where m and n are positive integers and $m > n$. Then there exists a unit p in $\mathcal{M}_m(D)$ such that the last $(m - n)$ rows of pa are all zero.*

Proof. The result is trivial if $a = 0$. Suppose that $a \neq 0$. Suppose that $n = 1$. Then

$$a = \begin{bmatrix} a_{11} \\ \vdots \\ a_{m1} \end{bmatrix}.$$

Now $a_{i1} \neq 0$ for some $1 \leq i \leq m$. If $i = 1$ let $p_1 = a_{11}^{-1}e_{11} + \sum_{k \neq 1} e_{kk}$; otherwise let

$$p_1 = e_{i1} + a_{i1}^{-1}e_{1i} + \sum_{k \neq i,1} e_{kk} \in \mathcal{M}_m(D).$$

Then p_1 is a unit in $\mathcal{M}_m(D)$ and $p_1 a$ has first entry 1. Thus without loss of generality $a_{11} = 1$. Now let

$$p = \begin{bmatrix} 1 & 0 & 0 & 0 & \cdots & 0 \\ -a_{21} & 1 & 0 & 0 & \cdots & 0 \\ -a_{31} & 0 & 1 & 0 & \cdots & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdots & \cdot \\ -a_{m1} & 0 & 0 & 0 & \cdots & 1 \end{bmatrix} \in \mathcal{M}_m(D).$$

Then p is a unit in $\mathcal{M}_m(D)$ with inverse

$$p^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 & \cdots & 0 \\ a_{21} & 1 & 0 & 0 & \cdots & 0 \\ a_{31} & 0 & 1 & 0 & \cdots & 0 \\ \cdot & \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdots & \cdot \\ a_{m1} & 0 & 0 & 0 & \cdots & 1 \end{bmatrix}.$$

Moreover,

$$pa = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

This proves the result when $n = 1$.

Now suppose that $n \geq 2$. Let a_1 be the $m \times (n-1)$ matrix over D and let b be the $m \times 1$ matrix over D such that $a = [a_1 | b]$ (in the obvious notation). By induction there exists a unit q_1 in $\mathcal{M}_m(D)$ such that $q_1 a_1$ has last $m - (n-1)$ rows zero. It follows that

$$q_1 a = [q_1 a_1 | q_1 b] = \begin{bmatrix} c & d \\ e & f \end{bmatrix},$$

where c, d, e, f are, respectively, an $(n-1) \times (n-1)$ matrix, an $(n-1) \times$ matrix, the zero $(m-(n-1)) \times (n-1)$ matrix and an $(m-(n-1)) \times 1$ matrix over D . If $f = 0$ then the result is proved. If $f \neq 0$ then we can argue as in the case $n = 1$ to produce a unit q_2 in $\mathcal{M}_m(D)$ such that

$$q_2 q_1 a = \begin{bmatrix} c & d \\ e & g \end{bmatrix}$$

where g is the $(m-(n-1)) \times 1$ matrix with first entry 1 and all other entries zero. Thus, if $p = q_2 q_1$ then p is a unit in $\mathcal{M}_m(D)$ and the last $(m-n)$ rows of pa are zero, as required. \square

The proof of the next result is quite elementary. Recall that if R is a left Ore domain with left quotient division ring D then any element in D can be written in the form $c^{-1}r$ where $r \in R$, $0 \neq c \in R$. It is well known that if n is a positive integer and $q_i \in D$ ($1 \leq i \leq n$) then there exist $r_i \in R$ ($1 \leq i \leq n$), $0 \neq d \in R$ such that $q_i = d^{-1}r_i$ ($1 \leq i \leq n$). This gives the following result.

Lemma 2.3. *Let R be a left Ore domain with left quotient division ring D and let m be a positive integer. Let p be any unit in $\mathcal{M}_m(D)$. Then there exists a nonzero element c in R such that $cp \in \mathcal{M}_m(R)$.*

In the next result we return to the situation considered in Section 1. Let R be any ring and let m and n be positive integers. Let S denote the ring $\mathcal{M}_n(R)$ and let α be the mapping from the collection of n -generated submodules of the free right R -module $F = R_R^{(m)}$ to the collection of cyclic S -submodules of $\mathcal{M}_{m \times n}(R)$, as defined in Section 1.

Lemma 2.4. *With the above notation, let $N \subseteq L$ be n -generated R -submodules of F such that N is an essential submodule of L . Then $\alpha(N)$ is an essential S -submodule of $\alpha(L)$.*

Proof. Let $L = (b_{11}, \dots, b_{m1})R + \dots + (b_{1n}, \dots, b_{mn})R$, and let (b_{ij}) be the corresponding matrix in $\mathcal{M}_{m \times n}(R)$. Let $s = (c_{ij}) \in S$, where $c_{ij} \in R$ ($1 \leq i, j \leq n$), such that $(b_{ij})s \neq 0$. There exists $1 \leq k \leq n$ such that the k th column of $(b_{ij})s$ is not zero. Thus

$$0 \neq x = (b_{11}, \dots, b_{m1})c_{1k} + \dots + (b_{1n}, \dots, b_{mn})c_{nk} \in L.$$

There exists $r \in R$ such that $0 \neq xr \in N$. Let $t = re_{kk} \in S$. Then $0 \neq (b_{ij})st \in \alpha(N)$. It follows that $\alpha(N)$ is essential in $\alpha(L)$. \square

We shall require the following special case of Lemma 2.4.

Corollary 2.5. *With the above notation, let R be a semiprime right Goldie ring. Let $N \subseteq L$ be n -generated R -submodules of F such that N is an essential submodule of L . Let a be any nonzero element of $\alpha(L)$. Then there exists a regular element c in R such that $ac^* \in \alpha(N)$, where c^* is the diagonal matrix in S with all diagonal entries c .*

Proof. Let $a \in (b_{ij})S$ (in the above notation). Let a_k ($1 \leq k \leq n$) denote the columns of a . The proof of Lemma 2.4 shows that for each $1 \leq k \leq n$ there exists an essential right ideal E_k of R with

$$[0 \ \dots \ 0 \ a_k \ 0 \ \dots \ 0](E_k e_{kk}) \subseteq \alpha(N).$$

Let $E = E_1 \cap \dots \cap E_n$. Then E is an essential right ideal of R and hence E contains a regular element c of R [9, 2.3.4 and 2.3.5]. Now

$$ac^* = [a_1 \ \dots \ a_n]c^* \in \alpha(N). \quad \square$$

Proof of Theorem 2.1. Let R be a left and right Ore domain with quotient division ring D . Let n be a positive integer such that the free right R -module $R_R^{(n)}$ satisfies n -acc. To prove that every free right R -module satisfies n -acc it is sufficient to prove that every finitely generated free right R -module satisfies n -acc (see, for example, [3, Theorem 1.5]).

Let m be any positive integer. Let $F = R_R^{(m)}$. If $m \leq n$ then F satisfies n -acc. Suppose that $m \geq n+1$. Let $N_1 \subseteq N_2 \subseteq N_3 \subseteq \dots$ be any ascending chain of n -generated submodules of F . By [3, Lemma 1.1], for each $i \geq 1$, N_i has uniform dimension at most n . Thus, without loss of generality we can suppose that N_1 is essential in N_i for all $i \geq 1$.

By Corollary 1.2, $\alpha(N_1) \subseteq \alpha(N_2) \subseteq \alpha(N_3) \subseteq \dots$ is an ascending chain of cyclic S -submodules of $\mathcal{M}_{m \times n}(R)$, where $S = \mathcal{M}_n(R)$. For each $i \geq 1$, let $a_i \in \mathcal{M}_{m \times n}(R)$ such that $\alpha(N_i) = a_i S$. By Lemmas 2.2 and 2.3 there exist a unit p in $\mathcal{M}_m(D)$ and a nonzero element c in R such that $cp \in \mathcal{M}_m(R)$ and cpa_1 has its last $(m-n)$ rows all zero. By Corollary 2.5, for each $i \geq 1$, there exists a nonzero element d_i in R such that $cpa_i d_i^* \in cpa_1 S$. Thus the last $(m-n)$ rows of $cpa_i d_i^*$ are zero and hence the last $(m-n)$ rows of cpa_i are zero.

Consider the ascending chain $cpa_1 S \subseteq cpa_2 S \subseteq cpa_3 S \subseteq \dots$ in $\mathcal{M}_{m \times n}(R)$. By Corollary 1.3 $\beta(cpa_1 S) \subseteq \beta(cpa_2 S) \subseteq \beta(cpa_3 S) \subseteq \dots$ is an ascending chain of n -generated submodules of F . Moreover, for each $i \geq 1$, $\beta(cpa_i S)$ is contained in the submodule G of F consisting of all elements of F of the form $(r_1, \dots, r_n, 0, \dots, 0)$, where $r_i \in R$ ($1 \leq i \leq n$). But $G \cong R_R^{(n)}$ and hence, by hypothesis, G satisfies n -acc. Thus there exists a positive integer k such that $\beta(cpa_k S) = \beta(cpa_{k+1} S) = \beta(cpa_{k+2} S) = \dots$. By Lemma 1.4, if we now apply α we have $cpa_k S = cpa_{k+1} S = cpa_{k+2} S = \dots$. Now using the fact that $c \neq 0$ and p is a unit, we have $a_k S = a_{k+1} S = a_{k+2} S = \dots$. Finally applying β we obtain $N_k = N_{k+1} = N_{k+2} = \dots$. It follows that F satisfies n -acc. \square

If in Theorem 2.1 the ring R is commutative we can do rather better, as the next result shows. If $a \in \mathcal{M}_n(R)$, for any commutative ring R and positive integer n , then $\det(a)$ will denote the determinant of a .

Theorem 2.6. Let R be a commutative domain and let n be a positive integer such that the free R -module $R_R^{(n-1)}$ satisfies n -acc. Then every free R -module satisfies n -acc.

Proof. In view of Theorem 2.1 it is sufficient to prove that the free R -module $F = R_R^{(n)}$ satisfies n -acc. Let $S = \mathcal{M}_n(R)$ and let D denote the field of fractions of R . By Corollary 1.6, it is sufficient to prove that S satisfies right 1-acc. Let $a_1S \subseteq a_2S \subseteq a_3S \subseteq \cdots$ be any ascending chain of nonzero principal right ideals of S . By the proof of Theorem 2.1, we can suppose without loss of generality that a_1 has rank n , for otherwise there exists a unit p in $\mathcal{M}_n(D)$ and a nonzero element c in R such that cpa_i has zero last row for all $i \geq 1$. Now $\det(a_1)R \subseteq \det(a_2)R \subseteq \det(a_3)R \subseteq \cdots$, so there exists a positive integer k such that $\det(a_k)R = \det(a_{k+1})R = \det(a_{k+2})R = \cdots$.

Note that for all $i \geq k$, $a_k = a_i b_i$ for some $b_i \in \mathcal{M}_n(R)$ and $\det(a_k)R = \det(a_i)R$. Since $\det(a_k) = \det(a_i) \det(b_i) \neq 0$, it follows that $\det(b_i)$ is a unit in R and hence b_i is a unit in $S = \mathcal{M}_n(R)$. Thus $a_k S = a_{k+1} S = a_{k+2} S = \cdots$. It follows that F satisfies n -acc, as required. \square

Nicolas [10, Proposition 1.4] proved that if R is a commutative domain which satisfies 1-acc then every free R -module satisfies 1-acc. Now Theorem 2.6 gives at once:

Corollary 2.7. *Let R be a commutative domain which satisfies 2-acc. Then every free R -module satisfies 2-acc.*

3. Rings whose free modules have pan -acc

In this section, our concern is to give, in the spirit of [2, 5], a range of examples of rings whose (finitely generated) free modules satisfy n -acc, for some positive integer n , or pan -acc. As noted earlier, Heinzer and Lantz [7] give examples, for each positive integer n , of a commutative ring R_n which satisfies n -acc but not $(n+1)$ -acc, and hence not pan -acc.

Proposition 3.1. *Let R be a subring of a ring S and let A be an ideal of R such that A is a left ideal of S and the ring R/A is right perfect. Suppose further that there exists a positive integer n such that every (finitely generated) free right S -module satisfies n -acc. Then every (finitely generated) free right R -module satisfies n -acc.*

Proof. Let n be any positive integer. Let I be any nonempty index set and let $N_1 \subseteq N_2 \subseteq N_3 \subseteq \cdots$ be any ascending chain of n -generated submodules of the free right R -module $R_R^{(I)}$. In a natural way we can think of $R_R^{(I)}$ as an R -submodule of the right S -module $F = S_S^{(I)}$.

Clearly $N_1 S \subseteq N_2 S \subseteq N_3 S \subseteq \cdots$ is an ascending chain of n -generated S -submodules of F . By hypothesis, there exists a positive integer t such that $N_t S = N_{t+1} S = N_{t+2} S = \cdots$. Because A is a left ideal of S it follows that $N_t A = N_{t+1} A = N_{t+2} A = \cdots$.

Let $N = \bigcup_{i \geq 1} N_i$. Then $NA = N_t A$ and hence N/N_t is a right (R/A) -module. By the Jonah–Renault Theorem (see [8, Main Theorem; 11, Proposition 1.2]), N/N_t satisfies n -acc and hence there exists $s \geq t$ with $N_s = N_{s+1} = N_{s+2} = \cdots$. \square

Now suppose that in Proposition 3.1, A is a finitely generated right ideal, rather than a left ideal, of S and that A_S is generated by k elements. In this case, in the proof of Proposition 3.1, $N_1A \subseteq N_2A \subseteq N_3A \subseteq \cdots$ is an ascending chain of (nk) -generated S -submodules of F . If F satisfies (nk) -acc then there exists a positive integer t such that $N_tA = N_{t+1}A = N_{t+2}A = \cdots$. By the proof of Proposition 3.1, it follows that $R_R^{(t)}$ satisfies n -acc. We have thus proved the following companion to Proposition 3.1.

Proposition 3.2. *Let R be a subring of a ring S and let A be an ideal of R such that A is a finitely generated right ideal of S and the ring R/A is right perfect. Suppose further that every (finitely generated) free right S -module satisfies pan-acc. Then every (finitely generated) free right R -module satisfies pan-acc.*

Proposition 3.3. *Let T be a subring of a ring S and let B and C be ideals of T such that the rings T/B and T/C are right perfect and C is a finitely generated right ideal of T . Let R_1 and R_2 denote the subrings $T + SB$ and $T + CS$ of S , respectively.*

(i) *If n is a positive integer such that every (finitely generated) free right S -module satisfies n -acc then so too does every (finitely generated) free right R_1 -module.*

(ii) *If every (finitely generated) free right S -module satisfies pan-acc then so too does every (finitely generated) free right R_2 -module.*

Proof. (i) Note that SB is a left ideal of S and a two-sided ideal of R_1 such that $R_1/SB \cong T/(T \cap SB)$ which is right perfect, being a homomorphic image of T/B . Apply Proposition 3.1 to obtain that every free right R_1 -module satisfies n -acc.

(ii) Similar to (i). \square

Proposition 3.4. *Let S be any ring and let n be a positive integer such that every (finitely generated) free right S -module satisfies n -acc. Let T be a subring of S and let B be an ideal of T such that the ring T/B is right perfect. Let L be any left ideal of S such that $B + LB$ is a left ideal of S and let $R = T + LB$. Then every (finitely generated) free right R -module satisfies n -acc.*

Proof. Let $A = B + LB$. Then A is a left ideal of S and a two-sided ideal of R such that $R/A = (T + LB)/(B + LB) \cong T/(B + (T \cap LB))$ which is right perfect, being a homomorphic image of T/B . Apply Proposition 3.1. \square

Corollary 3.5. *Let S be any ring and let n be a positive integer such that every (finitely generated) free right S -module satisfies n -acc. Let T be a subring of S and let B be an ideal of T such that the ring T/B is right perfect. Let L be any left ideal of S such that $S = T + L$ and let $R = T + LB$. Then every (finitely generated) free right R -module satisfies n -acc.*

Proof. Because $S = T + L$, $B + LB = SB$ is a left ideal of S . Apply Proposition 3.4. \square

The next result is a companion to Corollary 3.5.

Proposition 3.6. *Let S be any ring such that every (finitely generated) free right S -module satisfies pan-acc. Let T be a subring of S and let B be an ideal of T such that B is finitely generated as a right ideal and the ring T/B is right perfect. Let E be any right ideal of S such that $S = T + E$ and let $R = T + BE$. Then every (finitely generated) free right R -module satisfies pan-acc.*

Proof. There exist a positive integer m and elements $b_i \in B$ such that $B = b_1T + \cdots + b_mT$. Now $B + BE = BS = b_1S + \cdots + b_mS$. Now apply Proposition 3.2. \square

Let S be a ring and let A be a right ideal of S . Then we define $\mathcal{I}(A) = \{s \in S : sA \subseteq A\}$. Then $\mathcal{I}(A)$ is the biggest subring of S in which A is a two-sided ideal and $\mathcal{I}(A)$ is called the *idealizer* of A in S . If A is a left ideal we can construct the idealizer $\mathcal{I}(A)$ in a similar way.

Proposition 3.7. *Let A be a left or right ideal of a ring S , let T be a right perfect subring of $\mathcal{I}(A)$ and let $R = T + A$.*

(i) *If A is a left ideal and n is a positive integer such that every (finitely generated) free right S -module satisfies n -acc then so too does every (finitely generated) free right R -module.*

(ii) *If A is a finitely generated right ideal and every (finitely generated) free right S -module satisfies pan-acc then so too does every (finitely generated) free right R -module.*

Proof. (i) By Proposition 3.1 since $R/A \cong T/(T \cap A)$ which is right perfect.

(ii) Similar to (i). \square

Corollary 3.8. *Let T be a right perfect subring of a ring S , let A be an ideal of S and let $R = T + A$. If n is a positive integer such that every (finitely generated) free right S -module satisfies n -acc then so too does every (finitely generated) free right R -module.*

Proof. By Proposition 3.7, for in this case $S = \mathcal{I}(A)$. \square

We next mention an interesting special case of Proposition 3.7.

Proposition 3.9. *Let A be a left or right ideal of a ring S such that the S -module S/A has finite composition length. Let $R = \mathcal{I}(A)$.*

(i) *If A is a left ideal and n is a positive integer such that every (finitely generated) free right S -module satisfies n -acc then so too does every (finitely generated) free right R -module.*

(ii) *If A is a finitely generated right ideal and every (finitely generated) free right S -module satisfies pan-acc then so too does every (finitely generated) free right R -module.*

Proof. The ring R/A is isomorphic to the endomorphism ring of the S -module S/A and hence R/A is semiprimary, whence right perfect, by [1, 28.8 and 29.3]. Apply Propositions 3.1 and 3.2. \square

Now we introduce some matrix examples. First, we prove the following result.

Proposition 3.10. *Let A and B be ideals of a ring R such that $AB = 0$, the ring R/B is right perfect and every (finitely generated) free right (R/A) -module satisfies n -acc, for some fixed positive integer n . Then every (finitely generated) free right R -module satisfies n -acc.*

Proof. Let F be a (finitely generated) free right R -module and let $N_1 \subseteq N_2 \subseteq N_3 \subseteq \dots$ be any ascending chain of n -generated submodules of F . Then $(N_1 + FA)/FA \subseteq (N_2 + FA)/FA \subseteq (N_3 + FA)/FA \subseteq \dots$ is an ascending chain of n -generated submodules of the (finitely generated) free right (R/A) -module F/FA . By hypothesis, there exists a positive integer k such that $N_k + FA = N_{k+1} + FA = N_{k+2} + FA = \dots$. Now $AB = 0$ gives $N_k B = N_{k+1} B = N_{k+2} B = \dots$. The argument of Proposition 3.1 now gives that $N_t = N_{t+1} = N_{t+2} = \dots$ for some integer $t \geq k$. Thus F satisfies n -acc. \square

Let S and T be rings and let M be a left S -, right T -bimodule. Let $[S, M; 0, T]$ denote the set of “matrices”

$$\begin{bmatrix} s & m \\ 0 & t \end{bmatrix},$$

where $s \in S$, $m \in M$ and $t \in T$. Denote the above matrix by $[s, m; 0, t]$. Then $[S, M; 0, T]$ is a ring with respect to the usual definitions of matrix addition and multiplication.

Corollary 3.11. *Let S be a ring such that every (finitely generated) free right S -module satisfies n -acc, for some fixed positive integer n . Let T be a right perfect ring and let M be a left S -, right T -bimodule. Let $R = [S, M; 0, T]$. Then every (finitely generated) free right R -module satisfies n -acc.*

Proof. Let $A = [0, M; 0, T]$ and $B = [S, M; 0, 0]$. Then A and B are ideals of R , $AB = 0$, $R/A \cong S$ and $R/B \cong T$. Apply Proposition 3.10. \square

Corollary 3.11 has the following immediate consequence.

Corollary 3.12. *Let K be a field and let S be a K -algebra such that every (finitely generated) free right S -module satisfies n -acc, for some fixed positive integer n . Let M be any left S -module and let $R = [S, M; 0, K]$. Then every (finitely generated) free right R -module satisfies n -acc.*

Corollary 3.13. *Let K be a field and let S be a right and left Noetherian K -algebra. Let M be any left S -module and let $R = [S, M; 0, K]$. Then every free right R -module satisfies pan-acc.*

Proof. By Corollary 3.12 and [11, Corollaire 3.3]. \square

Example 3.14. Let K be a field and let S be a simple right and left Noetherian K -algebra which is not Artinian, let U be any simple left S -module and let $R = [S, U; 0, K]$. Then

- (i) R is a left Noetherian ring, every finitely generated free left R -module is Noetherian but not every free left R -module satisfies *l-acc*.
- (ii) Every free right R -module satisfies *pan-acc*.

Proof. (i) By [9, 1.1.7] and [11, Proposition 3.4].

(ii) By Corollary 3.13. \square

Now taking K, S as in Example 3.14 and U a simple right S -module, let R denote the ring $[K, U; 0, S]$. Let $A = [0, U; 0, S]$ and $B = [K, U; 0, 0]$. Then A and B are ideals of R and $AB = 0$. Moreover, $R/A \cong K$, so that R/A is right perfect, $R/B \cong S$, so that every free right (R/B) -module satisfies *pan-acc* [11, Corollaire 3.3], but not every free right R -module satisfies *l-acc* [11, Proposition 3.4]. Compare Proposition 3.10. Note also that in this case if $C = BA = [0, U; 0, 0] \neq 0$, then $C^2 = 0$ and $R/C \cong S \oplus K$, so that every free right (or left) (R/C) -module satisfies *pan-acc*.

Many more examples can be produced using Corollary 3.11. For example, let S be a commutative Noetherian domain with field of fractions L , let K be any extension field of L and let V be any vector space over K . Then the ring $R = [S, V; 0, K]$ has the property that every free right R -module satisfies *pan-acc* (Corollary 3.11 and [11, Corollaire 2.3]). Note that R is right Noetherian if and only if R is right Goldie if and only if V is finite dimensional over K [9, 1.1.7].

Our next aim is to give an example of a commutative domain R such that every free R -module satisfies *pan-acc* but the polynomial ring $R[t]$ does not satisfy *2-acc*. In contrast we have the following elementary fact.

Proposition 3.15. *Let R be a domain which satisfies right *l-acc*. Then the polynomial ring $R[t]$ satisfies right *l-acc*.*

Proof. Let S denote the ring $R[t]$. For any polynomial $f(t)$ in S , let $\delta(f(t))$ denote the degree of $f(t)$ and, if $f(t) \neq 0$, let $\lambda(f(t))$ denote the leading coefficient of $f(t)$.

Let $f_1(t)S \subseteq f_2(t)S \subseteq f_3(t)S \subseteq \dots$ be any ascending chain of principal right ideals of S . Then $\delta(f_1(t)) \geq \delta(f_2(t)) \geq \delta(f_3(t)) \geq \dots$, so that without loss of generality we can suppose that all the polynomials $f_i(t)$ are nonzero with the same degree.

Moreover, $\lambda(f_1(t))R \subseteq \lambda(f_2(t))R \subseteq \lambda(f_3(t))R \subseteq \dots$ so there exists a positive integer n with $\lambda(f_n(t))R = \lambda(f_{n+1}(t))R = \lambda(f_{n+2}(t))R = \dots$. It is now easy to check that $f_n(t)S = f_{n+1}(t)S = f_{n+2}(t)S = \dots$. Thus S satisfies right *l-acc*. \square

Example 3.16. Let K/L be a nonalgebraic field extension and let R denote the subring $L + xK[x]$ of the polynomial ring $K[x]$. Then R is a commutative domain such

that every free R -module satisfies *pan-acc* but the polynomial ring $R[t]$ does not satisfy *2-acc*.

Proof. Let T denote the ring $R[t]$. Note first that every free R -module satisfies *pan-acc* by Corollary 3.8 (take $S = K[x]$, $T = L$ and $A = xK[x]$). There exists an element a in K such that a is not algebraic over L . For each positive integer n ,

$$x^2 a^n = (x^2 a^{n+1})t - (xat - x)xa^n \in (x^2 a^{n+1}, xat - x).$$

Consider the chain of 2-generated ideals of T :

$$(x^2 a, xat - x) \subseteq (x^2 a^2, xat - x) \subseteq (x^2 a^3, xat - x) \subseteq \dots \quad (1)$$

Now suppose that $x^2 a^{n+1} \in (x^2 a^n, xat - x)$, for some positive integer n . There exist u, v in T such that

$$x^2 a^{n+1} = x^2 a^n u + (xat - x)v.$$

Setting $t = 1/a$, we have

$$x^2 a^{n+1} = x^2 a^n (d_0 + d_1(1/a) + d_2(1/a)^2 + \dots + d_m(1/a)^m)$$

for some $m \geq 1$, $d_i \in R$ ($0 \leq i \leq m$). Hence

$$a^{m+1} = d_0 a^m + d_1 a^{m-1} + \dots + d_m.$$

For each $0 \leq i \leq m$, there exist $c_i \in L$, $f_i(x) \in K[x]$ such that $d_i = c_i + x f_i(x)$. It follows that $a^{m+1} = c_0 a^m + c_1 a^{m-1} + \dots + c_m$, a contradiction. Thus every inclusion in the chain (1) is proper and hence the ring T does not satisfy *2-acc*. \square

Note that in Example 3.16 the ring $R[t]$ is isomorphic to the subring $L[t] + xK[x, t]$ of the polynomial ring $S = K[x, t]$. The ring S is a commutative Noetherian domain and every free S -module satisfies *pan-acc*. [11, Corollaire 2.3]. Moreover, the ring $R[t]$ has as a subring the ring $S' = L + xK[x, t]$. By Corollary 3.8 every free S' -module satisfies *pan-acc*. This indicates how vital it is to have a right perfect subring involved in the constructions in this section.

4. Torsionless modules

Let R be a ring and let M be a right R -module. The module M is called *torsionless* provided for each $0 \neq m \in M$ there exists $f \in \text{Hom}_R(M, R)$ such that $f(m) \neq 0$ (see, for example, [9, 3.4.2]). It is easy to see that this is equivalent to saying that M embeds in a direct product of copies of R_R . Note that if U is any left R -module then the right R -module $\text{Hom}_R(U, R)$ is torsionless (see [9, 3.4.2]).

Let R be a right Noetherian right nonsingular ring. Then every torsionless right R -module satisfies *pan-acc* (see [3, Theorem 1.5] or [11, Corollaire 2.3]). In this section

we shall give some examples of rings which need not be right Noetherian but for which every torsionless right module satisfies *pan-acc*.

Proposition 4.1. *Let R be a subring of a ring S and let A be an ideal of R such that A is a left ideal of S and the ring R/A is right perfect. Suppose further that there exists a positive integer n such that every torsionless right S -module satisfies n -acc. Then every torsionless right R -module satisfies n -acc.*

Proof. By the proof of Proposition 3.1 with the direct product $(R_R)^I$ replacing the direct sum $(R_R)^{(I)}$. \square

In a similar way, the proof of Proposition 3.2 can be adapted to give:

Proposition 4.2. *Let R be a subring of a ring S and let A be an ideal of R such that A is a finitely generated right ideal of S and the ring R/A is right perfect. Suppose further that every torsionless right S -module satisfies *pan-acc*. Then every torsionless right R -module satisfies *pan-acc*.*

Corollary 4.3. *Let R be a subring of a right Noetherian right nonsingular ring S and let A be an ideal of R such that A is a left or right ideal of S and the ring R/A is right perfect. Then every torsionless right R -module satisfies *pan-acc*.*

Proof. By Propositions 4.1 and 4.2 and [11, Corollaire 2.3]. \square

Another consequence of Propositions 4.1 and 4.2 is the following result.

Corollary 4.4. *Let T be a right Noetherian right nonsingular ring and let B be any ideal of T such that the ring T/B is right Artinian. Let R denote the subring $T + xB[x]$ of the polynomial ring $T[x]$. Then every torsionless right R -module satisfies *pan-acc*.*

Proof. If E is an essential right ideal of the polynomial ring $S = T[x]$ then the set E' of leading coefficients of the elements of E , together with 0, forms an essential right ideal of T . It follows that the ring S is right Noetherian right nonsingular. Let L denote the ideal xS of S . Note that $S = T + L$. Let $A = B + xB[x] = SB$. Then $A \subseteq R$ and A is an ideal of S . Moreover, the ring R/A is a homomorphic image of T/B , so is right Artinian. By Corollary 4.4, every torsionless right R -module satisfies *pan-acc*. \square

It is now clear that the results of Section 3 can be adapted to give corresponding results for torsionless modules. We now prove an analogue of Proposition 3.10.

Proposition 4.5. *Let A and B be ideals of a ring R such that $AB = 0$, the ring R/B is right perfect and every torsionless right (R/A) -module satisfies n -acc, for some fixed positive integer n . Then every torsionless right R -module satisfies n -acc.*

Proof. Let $F = (R_R)^I$, for any nonempty index set I . Let $R' = R/A$ and $A^* = A^I$. Then $F/A^* \cong (R'_R)^I$, which is a torsionless right R' -module. Now the result follows by the proof of Proposition 3.10 because $A^*B = 0$. \square

Corollary 4.6. *Let K be a field and let S be a right Noetherian right nonsingular K -algebra. Let M be any left S -module and let $R = [S, M; 0, K]$. Then every torsionless right R -module satisfies pan-acc.*

Proof. This result follows from Proposition 4.5 in essentially the same way that Corollary 3.12 follows from Proposition 3.10, by using [11, Corollaire 2.3]. \square

Acknowledgements

This work was carried out during the visits of the first author to the University of Glasgow in December 1994 and February 1995, and she would like to thank the Department of Mathematics of the University of Glasgow for hospitality and the JNICT (Portugal) for financial support.

References

- [1] F.W. Anderson and K.R. Fuller, Rings and Categories of Modules (Springer, New York, 1974).
- [2] M.E. Antunes Simões and A.-M. Nicolas, Exemples d'anneaux n -acc, Comm. Algebra 12 (1984) 1653–1665.
- [3] M.E. Antunes Simões and P.F. Smith, Direct products satisfying the ascending chain condition for submodules with a bounded number of generators, Comm. Algebra 23 (1995) 3525–3540.
- [4] M.E. Antunes Simões and P.F. Smith, On the ascending chain condition for submodules with a bounded number of generators, Comm. Algebra 24 (1996) 1713–1721.
- [5] F. Bonang, Noetherian rings whose subidealizer subrings have pan-a.c.c., Comm. Algebra 17 (1989) 1137–1146.
- [6] P.M. Cohn, Free Rings and Their Relations (Academic Press, New York, 1971).
- [7] W. Heinzer and D. Lantz, Commutative rings with ACC on n -generated ideals, J. Algebra 80 (1983) 261–278.
- [8] D. Jonah, Rings with the minimum condition for principal right ideals have the maximum condition for principal left ideals, Math. Z. 113 (1970) 106–112.
- [9] J.C. McConnell and J.C. Robson, Noncommutative Noetherian Rings (Wiley-Interscience, Chichester, 1987).
- [10] A.-M. Nicolas, Sur les modules tels que toute suite croissante de sous-modules engendrés par n générateurs soit stationnaire, J. Algebra 60 (1979) 249–260.
- [11] G. Renault, Sur les conditions de chaînes ascendantes dans des modules libres, J. Algebra 47 (1977) 268–275.